



Emergency Medication Management Plan

Midazolam (Zyamis Prefilled Syringe)

(only to be administered by a trained person)

Midazolam – Zyamis Management Plan for:

Date of Birth: text

Date of plan: text

Weight: text

Administration method: BUCCAL

1. FIRST DOSE Midazolam (Zyamis)

First dose prefilled syringe =

☐ 5mg/0.5ml (Blue) ☐ 7.5mg/0.75ml (Purple) ☐ 10mg/1ml (Orange)

For single seizures:

- ☐ As soon as a text (seizure type) begins
- ☐ If the text (seizure type) continues longer than text mins text hrs

Special instructions: text

For clusters of seizures:

- ☐ When text (number) text (seizure type) occurs within text mins text hrs
- ☐ Other (please specify): text

Special instructions: text

2. SECOND DOSE Midazolam (Zyamis)

Second dose prefilled syringe =

☐ 5mg/0.5ml (Blue) ☐ 7.5mg/0.75ml (Purple) ☐ 10mg/1ml (Orange)

- ☐ Not prescribed OR
- ☐ If the text (seizure type) continues for another text mins following the first dose
- ☐ When another text (number) text (seizure type) occur/s within text mins text hrs following the first dose
- ☐ Other (please specify): text

Special instructions: text

3. Maximum number of Midazolam doses to be given in a 24-hour period

Maximum number:

4. Dial 000 to call the ambulance:

- ☐ Prior to administering Midazolam
- ☐ Immediately after administering Midazolam
- ☐ If the seizure has not stopped minutes after giving the midazolam
- ☐ If injured or has difficulty with breathing
- ☐ Other (please specify):

5. Describe what to do after Midazolam has been administered:

- Continue to provide seizure first aid, keeping me safe from injury. Continue to time and monitor the seizure.
- If possible, keep me on my side to protect my airway. If this is not possible, turn my head to the side.
- If in a wheelchair or car seat, leave me seated and continue to support my head and protect my airway.
- Document time Midazolam was given, when ambulance called and when seizure activity stops.
- Stay with me and monitor:
 - For shallow or slow breathing, change of facial colour
 - Other:

6. Prescribing medical practitioner

Name of medical practitioner:

Telephone:

Date:

Signature:

7. Storage and family special instructions

Recommended MIDAZOLAM storage information:

- Keep out of reach of children
- Protect from light (stored in container provided) and store at room temperature (below 25°C)
- Regularly check the expiry date.

Any special instructions e.g. storage of medication, when on outings etc. or people to contact if emergency medication is given:

Emergency contact name: text	
Relationship: text	Telephone: text
Signature: text	Date: text

PO Box 1834, Launceston TAS 7250
31 Thistle St, South Launceston 7249
1-7 Liverpool St, Hobart TAS 7000
P: 6344 6881/1300 852 853
www.epilepsytasmania.org.au